

AIR FORCE INV. NO. AFD 489  
HARVEY A. SCHWERTNER ET AL.  
SHEET ONE OF SIX

CLINICAL AND LABORATORY CHARACTERISTICS OF 644 STUDY

SUBJECTS BY DEGREE OF CAD

CAD RISK FACTORS <sup>a</sup>	Maximum stenosis			P-value
	0 - 9% (n = 430)	10 - 49% (n = 98)	50 - 100% (n = 116)	
Age, years	39.9 (6.3)	45.3 (5.9)	45.3 (5.4)	.000
Cigarettes/day	5.68 (10.91)	7.84 (13.57)	10.59 (13.87)	.000
Systolic blood pressure, mm Hg	125.7 (13.2)	126.8 (13.0)	131.0 (13.3)	.001
Total bilirubin, $\mu$ mol/L	14.8 (7.1)	13.1 (4.7)	12.5 (5.8)	.001
Total cholesterol, mmol/L	5.32 (0.99)	5.91 (0.93)	6.16 (1.15)	.000
HDL-cholesterol, mmol/L	1.21 (0.31)	1.14 (0.26)	1.11 (0.24)	.002
Triglycerides <sup>b</sup> , mmol/L	1.56 (1.11)	1.76 (0.92)	1.87 (0.88)	.009
Cholesterol/HDL-C <sup>c</sup>	4.71 (1.62)	5.42 (1.47)	5.81 (1.59)	.000
(Cholesterol/bilirubin) ( $\div$ 100) <sup>d</sup>	4.26 (1.91)	5.11 (1.95)	5.85 (2.56)	.000
Cholesterol/(HDL-C + bilirubin) <sup>c</sup>	1.69 (0.59)	2.01 (0.57)	2.21 (0.66)	.000
LDL-C/HDL-C <sup>c</sup>	3.01 (1.11)	3.64 (1.16)	3.97 (1.33)	.000
LDL-C/(HDL-C + bilirubin) <sup>c</sup>	1.08 (0.42)	1.35 (0.43)	1.51 (0.53)	.000

<sup>a</sup> Values are given as mean  $\pm$  SD. To convert values for cholesterol, triglycerides, and bilirubin to mg/dL, multiply by 38.66, 88.54, and 0.05847. Analysis of variance. All F-values were significant at P = 0.009.

<sup>b</sup> Triglyceride concentrations are expressed as mmol/L triolein.

<sup>c</sup> Cholesterol/HDL-C, LDL-C/HDL-C, cholesterol/(HDL-C + bilirubin), and LDL-

C/(HDL-C + bilirubin) ratios are presented on a mg/dL basis. Bilirubin concentrations

were multiplied by 100 before combining with HDL-C. For example, the cholesterol/(HDL-C + bilirubin) ratio was calculated by taking each individual's cholesterol concentration in mg/dL and dividing it by the sum of the HDL-cholesterol concentration in mg/dL plus the bilirubin concentration in mg/dL  $\times$  100.

<sup>d</sup> Cholesterol/bilirubin ratios were calculated as the ratio of cholesterol, mmol/L, divided by bilirubin, mmol/L, divided by 100.

*FIG. 1*

AIR FORCE INV. NO. AFD 489  
HARVEY A. SCHWERTNER ET AL.  
SHEET TWO OF SIX

RESULTS OF DISCRIMINANT ANALYSIS USING TRADITIONAL  
RISK FACTORS AND VARIOUS LIPID-LIPOPROTEIN AND LIPID-  
LIPOPROTEIN-BILIRUBIN COMBINATIONS

Discriminant analysis showing variables tested for inclusion into model.

Variables accepted in model	Wilks' Lambda
Age	0.848
LDL-C/(HDL-C + bilirubin)	0.839
Systolic blood pressure	0.769
Variables not included in model	
Cholesterol/(HDL-C + bilirubin)	0.757
HDL-cholesterol	0.757
Triglycerides	0.757
Cholesterol/HDL-C	0.755
Cholesterol/bilirubin	0.755
LDL-C/HDL-C	0.753
Total bilirubin	0.751
Cigarettes per day	0.749
Total cholesterol	0.749

Coronary artery disease classification results.

Actual group	No. cases	Predicted Group Membership		
		0 - 9%	10 - 49%	50 - 100%
0 - 9%	457	430 94.1%	3 0.7%	24 5.3%
10 - 49%	98	74 75.5%	6 6.1%	18 18.4%
50 - 100%	116	74 63.8%	1 0.9%	41 35.3%

*FIG. 2*

AIR FORCE INV. NO. AFD 489  
HARVEY A. SCHWERTNER ET AL.  
SHEET THREE OF SIX

PERCENT CORRECT CLASSIFICATION FOR EACH RISK FACTOR  
COMBINATION FOR PREDICTING CORONARY ARTERY DISEASE

Coronary Artery Disease Classification <sup>a</sup>

Risk Factor Variable	0 - 9% Specificity %	10 - 49%	50 - 100% Sensitivity %	Overall Correct %
Age	97.2	0	13.8	67.4
Age, cholesterol	95.1	4.1	23.3	68.3
Cholesterol/HDL-C	97.0	0	6.8	67.2
LDL-C/HDL-C	97.5	0	6.9	67.6
Cholesterol/bilirubin	96.8	0	14.4	68.3
Cholesterol/(HDL-C + bilirubin)	96.1	0	17.1	68.3
LDL-C/(HDL-C + bilirubin)	96.0	0	17.9	68.4
Laboratory risk factors <sup>b</sup>	95.1	0	19.2	68.0
Traditional risk factors <sup>c</sup>	92.8	8.2	28.4	68.3
Traditional and cholesterol/(HDL-C + bilirubin) <sup>d</sup>	92.1	8.2	35.3	69.1
Traditional and LDL-C/(HDL-C + bilirubin) <sup>e</sup>	94.1	6.1	35.3	71.1

<sup>a</sup> 95% Confidence intervals for sensitivity, specificity, and overall correct are approximately  $\pm 8\%$ ,  $\pm 2\%$ , and  $\pm 3\%$ , respectively.

<sup>b</sup> Laboratory risk factors that entered the discriminant model were LDL/(HDL + bilirubin) and cholesterol.

<sup>c</sup> Traditional risk factors examined included cholesterol, cigarettes/day, systolic blood pressure, triglycerides, cholesterol/HDL-C, and HDL-cholesterol. Age, cholesterol, systolic blood pressure, and cigarettes/day entered the discriminant model.

<sup>d</sup> Age, cholesterol/(HDL-C + bilirubin), cholesterol, systolic blood pressure, and cigarettes/day entered the discriminant model.

<sup>e</sup> Age, LDL/(HDL-C + bilirubin), and systolic blood pressure entered the model.

*FIG. 3*

AIR FORCE INV. NO. AFD 489  
HARVEY A. SCHWERTNER ET AL.  
SHEET FOUR OF SIX

DIAGNOSTIC PERFORMANCE OF VARIOUS RISK FACTORS FOR  
PREDICTING SEVERE CORONARY ARTERY DISEASE USING 75TH  
PERCENTILE AS A CUT-POINT


Risk Factor Variable <sup>a,c</sup>	Sensitivity (%)	Specificity (%)	Efficiency of Test (%) <sup>b</sup>
Cholesterol/HDL-C	(59/146) 40.4	(573/731) 78.4	(632/877) 72.1
LDL-C/HDL-C	(63/145) 43.4	(577/731) 78.9	(640/877) 73.0
Cholesterol/bilirubin	(71/146) 48.6	(584/730) 79.9	(655/877) 74.7
Cholesterol/(HDL-C + bilirubin)	(76/146) 52.1	(588/731) 80.4	(664/877) 75.7
LDL-C/(HDL-C + bilirubin)	(75/145) 51.7	(587/731) 80.3	(662/877) 75.5

<sup>a</sup> Numbers in parentheses represent the actual subject counts. Cut-points (75th percentile) for the cholesterol/HDL-C ratio, LDL-C/HDL-C ratio, cholesterol/bilirubin ratio, cholesterol/(HDL-C + bilirubin) ratio, and LDL-C/(HDL-C + bilirubin) ratio were 6.04, 4.12, 3.84, 2.25, and 1.5, respectively.


<sup>b</sup> Efficiency of test was calculated as follows:  $TP + TN / (TP + TN + FP + FN)$

<sup>c</sup> 95% Confidence intervals for sensitivity, specificity, and efficiency of a test are approximately  $\pm 8\%$ ,  $\pm 4\%$ , and  $\pm 3\%$ , respectively.

*FIG. 4*



*FIG. 5*



*FIG. 6*